J. R. Statist. Soc. A (2008)
171, Part 1, pp. 137–157

The analysis of change, Newton's law of gravity and association models

Mark de Rooij

Leiden University, The Netherlands

[Received September 2004. Final revision April 2007]

Summary. Newton's law of gravity states that the force between two objects in the universe is proportional to the product of their masses and inversely proportional to the square of the distance between the two objects. In the first part of the paper this inverse-square model is cast in a general intergeneration gravities and to the analysis of longitudinal categorical data where the number of categories is the same for the two occasions. Symmetry, quasi-symmetry and Rasch type models and the goal is to obtain estimates of mass for the two categories and an estimate of the distance between the two categories. Two new models for the analysis of change in square mobility tables are introduced. It is shown that this generalized law of gravity is equivalent to Zeeman's model for social change and to the Bradley–Terry model when there are only two categories or only two kinds of first-stage data. The first case is when there are multiple two-way tables, and in the second case we have change over three points in time. Conditional and partial generalizations of the inverse-square law are also discussed. The models are illustrated by using Dutch voting data and models respectively.

Keywords: Categorical data; Euclidean distance; Gravity model; Longitudinal data; Square tables; Triadic distance

1. Introduction

This paper will be concerned with longitudinal *categorical data*, i.e. *repeated* measurements on a categorical variable with observational units with the same instrument. The main interest in studying longitudinal data is whether change occurred and, if so, what the nature of the change is. We shall confine ourselves to the case of categorical data. One questions concerns qualitative change, i.e. changes in attitude, opinion, behaviour or any other categorical variable. This is typically studied by investigating the structure of square tables. Another type of change is quantitative change or *mover*, for categorical data descriptions are in terms of 'different' or 'the same'.

Once longitudinal categorical data have been collected they can be represented in transition frequencies in a square table. Each cell in such a table contains the number of observations that have a variable measured at a specific time point (we adopt the way-mode terminology for the tables of Coombs and Kao (1955)). The number of time points defines the number of ways of the transition frequency table. Having measured a group of people twice on a categorical variable, a square transition frequency table arises. If measurements are obtained at three time points the data can be collected in a three-way square table, and so on.

An example of such data is obtained from Upton and Sinfield (1981) who discussed changes in political voting in Sweden. The data are shown in Table 1. There are five political parties: the Communists (COM), the *Social Democrats* (SD), the *Centre Party* (C), the *People's Party* (P; the

Address for correspondence: Mark de Rooij, Methodology and Statistics Unit, Institute for Psychological Research, Leiden University, PO Box 9555, 2300 RB Leiden, The Netherlands.
E-mail: mderooij@fsw.leidenuniv.nl

© 2008 Royal Statistical Society 0964-1998/08/171137

Table 1. *Swedish voting data representing voting changes from 1982 (rows) to 1979 (columns)*[1]

	COM	SD	C	P	VPK
COM	124	47	3	0	14
SD	7	1611	19	5	7
C	0	61	1022	19	0
P	0	26	63	1363	0
VPK	11	9	0	0	113

[1]From Upton and Sinfield (1981).

Conservatives CON). These are the anglicized names following Upton and Sinfield (1981). The rows correspond to the political parties in 1982 (in capital letters), the columns to the political parties in 1979 (in capital letters).

The focus will be on change, i.e. on the off-diagonal entries. The values on the diagonal are within-parties/movers, for these cells special parameters (which are often called loyalty parameters) will be introduced in the models that follow.

The question, once we have such change data, is not whether there is association but what the *pattern* of association looks like. We shall propose a model for these data based on Newton's law of gravity, which states that the force between any two objects in the universe is proportional to the masses of the two objects and inversely related to the squared distance between the two objects. Although this law is from physics, we shall show that it can be used in the social sciences. The dimensionals model will be applied to the analysis of change where the objects to be the universe are the categories of the variable. The force between two objects is measured by the number of observations in the off-diagonal cells of the square table. The mass of the objects is, however, unmeasured, however, since a sample is obtained from a population. Therefore, the law of gravity is used as a probabilistic model assuming a multinomial sampling scheme (which is the usual sampling scheme for square tables). The mass of the objects is explained by the mass of the two categories and a function of the distance between the two objects.

The remainder of this paper is organized as follows. The next section discusses Newton's law of gravity in more detail. Section 3 describes the analysis of change in terms of Newton's law of gravity and introduces dynamic elements in the law to adapt for different data settings. After that, in Section 4, the models are applied to the Swedish voting data. Some relationships with the RC(*M*) association model (Goodman, 1979, 1981). The usual identification constraints for this model, however, are not suited to the analysis of change. A new way of identifying the parameters is proposed in Section 5. Section 6 discusses the case of multiple two-way tables. The model will be generalized to the case of multiple two-way tables. The gravity models that are developed are related to conditional association models (Clogg, 1982), but again the usual identification constraints are not suited for the gravity models. Section 7 discusses the gravity models for three time points. These models are related to partial association models (Clogg, 1982). Identification and an application to empirical data will be discussed. We shall conclude with discussion and reflection about the results obtained and show some limitations of the work presented.

2. Newton's law of gravity

One of the major laws of the natural sciences is Newton's law of gravity:

> 'All matter attracts all other matter with a force proportional to the product of their masses and inversely proportional to the square of the distance between them'.

Fig. 1. Newton's law of gravity: the masses of objects x and y are represented by the axes of the circles and d_{xy} is the distance between the centres of the two objects

This law can be written in a formula as

$$F_{xy} = \frac{m_x m_y \alpha}{d_{xy}^2}, \tag{1}$$

where F_{xy} denotes the force between objects x and y, $m(x)$ and $m(y)$ are the masses of the objects x and y, and d_{xy}^2 is the squared distance between the objects. A more general formulation of the law is

$$F_{xy} = \frac{m_x m_y \alpha}{d_{xy}^\beta}, \tag{2}$$

where $\alpha / (\log(n)n^\beta)$ but may also be some other function. A graphical representation is shown in Fig. 1, where the masses are represented by the area of a circle. Newton explained a wide range of previously unrelated phenomena by using this law: the eccentric orbits of comets, the tides and their variations, the precession of the Earth's axis and motion of the Moon as perturbed by the pull of the Sun on it. Newton's law of gravity is considered to be universally valid.

In the next section we shall show that the law of gravity applies well to the analysis of social change. Therefore, first some other definitions of the function $g(\cdot)$ are provided and dynamic elements are added to the law. The dimensionals model will be applied to longitudinal data. It leaves model in statistics and social research, the RC(*M*) association model (Goodman, 1979, 1981). It should be noted, however, that by the time that we arrive at the RC(*M*) association model many properties of real forces as they are in the natural sciences have been lost. What is maintained is the interpretation in terms of mass and distance, and the analogy with Newton's law of gravity is meant more like a metaphor than reality.

3. The analysis of change

Where the task for Newton was to assess the force of the two objects on each other given their mass and their distance, we deal with the reverse problem. We assume that each object (category) has a mass and that the force between the objects is a function of the number of people between objects. These flows can be considered measurements of the attractional forces between objects, and then (using the law of gravity) are the number of people going from one object to another proportional to the mass of the first object times the mass of the second object and inversely proportional to a function of the distance between the two objects.

Table 1 is an illustrative example. The mass of the Communists and the Social Democrats in 1982 and for the Centre Party in 1979, or there is a large force between these two categories. Similarly, the force between the Communists and the Conservatives is small (apparently the Communists and the Conservatives are not attracted to each other). The reverse, for example, the force between Communists Social Democrats equals 47 and the force Social Democrats–Communists equals 31.

3.1. A Gaussian link

In Newton's law of gravity the distance is defined by a three-dimensional Euclidean distance, i.e. our universe is three dimensional. For the analysis of change the dimensionality is not known in advance. Therefore, we shall denote by S the number of dimensions. The objects will be the parties are differentiated on the standard 'left–right' dimension that is often found in political systems. Furthermore, there might be another dimension that differentiates the five parties. In this case $S = 2$. Of course, S can be greater than 2, but we shall restrict ourselves to the case $S = 2$.

Often the dimensions are interpreted after the solution has been found, on the basis of practical knowledge of the data at hand. The sth coordinate of object i in S-dimensional space are given by the vector $\mathbf{x}_i' = (x_{i1}, \ldots, x_{iS})$. The $n(n-1)$ rows, be collected in the $I \times S$ matrix $\mathbf{X} = (\mathbf{x}_1', \mathbf{x}_2', \ldots, \mathbf{x}_I')'$. The squared Euclidean distance between objects i and j is given by

$$d_{ij}^2(\mathbf{X}) = \sum_{s=1}^{S} (x_{is} - x_{js})^2. \tag{3}$$

Other distance measures might be used as well (e.g. any distance from the Minkowski family (see Borg and Groenen (2005), chapter 17)). Where in the law of gravity $g(d_{ij}) = d^{-2}$ we shall employ $g(\cdot) = \exp(-r^2)$, the Gaussian transformation or Gaussian link function (de Rooij and Heiser, 2005). Neperday, 1951), which is a monotone function. Again, like for the distance formulation, other transformation functions might be used, but in Section 7.3 it will be shown that this function makes the law of gravity to a well-known model for the statistical analysis of contingency tables.

3.2. Dynamic masses

As discussed above the measured forces are not symmetric, i.e. the force from Communist on Social Democrats is different from the force from Social Democrats on Communist is 16. The law of gravity assumes symmetric forces and the asymmetry is a form of 'error'.

To deal with such asymmetries the model will be generalized in two ways. The first generalization is to assume that the masses of the objects are dynamic, i.e. they change over time. The second is to extend the masses of the objects to be varied across m, i.e. object specific masses or a different point and unprovider of another. For example, in brand switching data some brands come into fashion at one moment and go out of fashion another. When an object is popular it has a large mass, when it is unpopular it has a small mass. For our model this means that objects have a mass at time point 1 and a mass at time point 2. Therefore, for object i we have m_{i1} at time point 1 for a graphical representation like Fig. 1), each object would have two circles.

3.3. Dynamic positions

A second generalization is to make the positions time dependent. So, *dynamic positions* are assumed. This is in line with Coleman's (1964) model for change. Coleman's model assumes that the object has changed. For example, a political party might change its election programme after it has lost dramatically in the last election or when a loss is to prospect, and thereby change its position on the political spectrum. Let $\mathbf{x}_{i1}' = (x_{i11}, \ldots, x_{i1S})$ and $\mathbf{x}_{i2}' = (x_{i21}, \ldots, x_{i2S})$ be denoted by $\mathbf{x}_i(1) = (x_{i11}, \ldots, x_{i1S})'$ and the positions of all objects at time point 1 are gathered in a matrix $\mathbf{Z}_1 = (\mathbf{x}_1(1), \mathbf{x}_2(1), \ldots, \mathbf{x}_I(1))'$. The one-mode Euclidean distance (3) is replaced by a two-mode Euclidean distance:

$$d_{ij}^2(\mathbf{Z}_1, \mathbf{Z}_2) = \sum_{s=1}^{S} (x_{i1s} - x_{j2s})^2. \tag{4}$$



Fig. 2. Graphical representation of Coleman's model: Swedish parties, N. COM, SD, C, P, VPK; mass and positions of the parties in 1982; α, size; p_s, mass and positions of the parties in 1979

1979; Becker, 1970; Huberman, 1993). By back-transforming using equations (7) and (5) we obtain the expected frequencies for the cells in table 1. Because of the large number of cells in the models, the transformation to a distance model and the way of identifying the solution are performed in MATLAB (Mathworks, 2004). The model with stable masses but dynamic positions is a special case of the symmetric association model and thus cannot be estimated with available software for association models.

3.4. Application to Swedish political data

The data in Table 1 were analysed by using the gravity models proposed. First some benchmark models are fitted. The independence model (corresponding to the model of complete homogeneity in models) do not fit these data (X^2 values respectively 321.13, 76.55 and 12.39); df 192, 12, 32 3 and 71.77, with 11, 10 and 6 degrees of freedom; df). Since the quasi-symmetry model does not fit the data, the symmetric association model (Goodman, 1979) does not fit the data either ($X^2 = 12.39$; $df = 6$). With one dimension $X^2 = 24.95$, $X^2 = 17.29$ and $df = 7$ and with two dimensions the fit barely increased: $X^2 = 12.05$, $G^2 = 11.25$ and $df = 3$. With dynamic positions a good fit was obtained in a single dimension, $X^2 = 4.76$, $G^2 = 5.57$ and $df = 6$. The solution is shown in Fig. 2. We see several positional changes there in 1982 the positions of the five parties are as expected on the left–right political spectrum. The masses in 1979 and 1982 are shown by the area of the circles. The Communists, the Social Democrats, the Centre Party, the People's Party and the Conservatives.

The positional changes from 1982 to 1979 can be summarized as follows: the Communists and Social Democrats grouped together on the left wing whereas the Conservatives and People's Party grouped on the right wing. Especially the Social Democrats made a big change to the left. The Centre Party stayed in the middle. The mass of the Communists in 1979 was 0. The mass of the Conservatives happened that distinguishes the two left-wing parties from the three other parties. Such a grouping was also found in Upton and Sinfield (1981). Although it may seem strange that the Social Democrats are more leftist than the Communists this has also been observed at several points in time by Lewis *et al.* (1972), page 228.

The mass of the Social Democrats in 1982 is the mass of the Communists, the People's Party and the Social Democrats stay the same, the Centre Party is the winner and the Conservative Party is the political party that lost mass.

4. Multiple two-way tables and multiple universes

Up to this point dimensions has been confined to two-way tables. In the remainder of this paper we shall generalize the models to three-way tables. This section treats the case of multiple

$$F_{xy} = \frac{m_x(\alpha m_y) / \exp(0.5\tau_x)}{\exp(\tau_x^2 / 2\sigma_x^2 \cdot \mathbf{Z}_1 \cdot \mathbf{Z}_2)}. \tag{6}$$

This gravity model can be rewritten in a form that is well known in statistics and is often applied in sociological studies; the RC(*M*) association model (Goodman, 1979, 1981). By back-transforming the parameters of this model, estimates of the masses and a co-ordinate of row gravities and column gravities are obtained. Identification constraints for the parameters of the RC(*M*) contingency tables are well established, and are then also valid for our gravity model. However, the usual graphical displays for the RC(*M*) association model are susceptible to misinterpretation (for examples see de Rooij and Heiser (2005)), where a wrong interpretation is more sensitive. The transformation from gravity to association model is as follows (de Rooij and Heiser, 2005):

$$\begin{aligned} F_{ij} &= \frac{m_i(\alpha m_j)}{\exp\left(\sum_{s=1}^{S} x_{is}^2 - 2x_{is}x_{js} + x_{js}^2\right)} \\ &= \frac{m_i(\alpha m_j)}{\exp\left(\sum_{s=1}^{S} x_{is}^2\right) \exp\left(\sum_{s=1}^{S} x_{is}x_{js}\right) \exp\left(\sum_{s=1}^{S} x_{js}^2\right)}. \end{aligned} \tag{5}$$

Defining $\alpha_i^{(1)} = m_i / \exp(\sum_{s=1}^{S} x_{is}^2)$ and $\beta_j^{(1)} = (\alpha m_j) / \exp(\sum_{s=1}^{S} x_{js}^2)$, we obtain

$$\begin{aligned} F_{ij} &= \frac{\alpha_i^{(1)} \beta_j^{(1)}}{\exp\left(\sum_{s=1}^{S} x_{is}x_{js}\right)} \\ &= \alpha_i \beta_j \exp\left(-\sum_{s=1}^{S} x_{is}x_{js}\right) \\ &= \alpha_i \beta_j \exp\left(-\sum_{s=1}^{S} \phi_s x_{is}x_{js}\right), \end{aligned} \tag{7}$$

with $\alpha_i = \alpha_i^{(1)}$, $\beta_j = \beta_j^{(1)}$ and $\phi_s = -1$. The last line is expression (7) in Goodman's (1979, 1991) RC(*M*) association model. In summary, we started with (an adaptation of) Newton's law of gravity, introduced dynamic elements and ended up with the well-known model. The RC(*M*) association model is a reduced rank model for the association which equals the number of dimensions S. The gravity model is a model for the mean, and the RC(*M*) association model is a mean model where the dimensionality is 0. The model with stable positions is the homogeneous RC(*M*) association model and imposes a symmetry restriction on the association, and thus is a quasi-symmetry model (Caussinus, 1965; Goodman, 1979, 1991). The model with stable masses and positions is a special case of the symmetry model.

Since our focus is on the off-diagonal entries we need parameters for the diagonal entries of the table. These are loyalty parameters for each class, i.e. the model becomes

$$F_{ij} = \begin{cases} \alpha_i \beta_j \exp\left(-\sum_{s=1}^{S} \phi_s x_{is}x_{js}\right) & \text{if } i \neq j, \\ \gamma_i \alpha_i \beta_j \exp\left(-\sum_{s=1}^{S} \phi_s x_{is}x_{js}\right) & \text{if } i = j, \end{cases} \tag{8}$$

where γ_i equals 1 if $i = j$ and 0 otherwise. The λ_i are object-specific loyalty parameters, of which there are I. The effect of these parameters is that the observations on the diagonal do not influence the masses and the distances, i.e. the gravity model persists to change. Another effect of these parameters is that the expected frequencies equal the observed frequencies for these cells.

3.5. Distances, distances and inner products

The RC(*M*) association model is often thought of as a model for ordinal data (although, strictly, it is a model for nominal categories). Since the RC(*M*) association model and gravity models are often interpreted in terms of distance. This raises the question what is now about the distance interpretation in the gravity model. To answer this question we should distinguish between the squared Euclidean distance and the inner product. The squared Euclidean distance between the set of row points can be interpreted such that, when $\mathbf{x}_i$ and $\mathbf{x}_{i'}$ with $\mathbf{x}_i = (x_{i1}, \ldots, x_{iS})'$ are (approximately) equal, categories i and i' have the same pattern of association to the column categories. Therefore, categories i and i' are classified in the same response category. The RC(*M*) association model can mean product role must be used, where the association is on the product of the length of the two vectors μ_i and ν_j multiplied by the cosine of the angle between these two vectors. The parameterization in terms of Newton's law of gravity provides a between-set distance interpretation, i.e. an interpretation of the distance between $\mathbf{x}_i(1)$ and $\mathbf{z}_j(2)$.

3.6. Identification constraints

The RC(*M*) association model is not identified; it needs location, scaling and cross-dimensional constraints on the row and column scores. Usually the scores are centred, the sum of squares is equal to 1 and the scores are uncorrelated (Goodman, 1979, 1991). Because of change, however, these standard identification constraints prevent substantial conclusions.

Let us denote the centred row scores by $\mathbf{x}_i(1)$. It is possible to transform these centred row scores to $\mathbf{x}_i(1) = \mathbf{T}^{-1}\mathbf{x}_i(1) + \mathbf{v}$, where $\mathbf{T}$ is a $S \times S$ matrix of rank S and $\mathbf{v}$ is a $S \times 1$ vector, and $\mathbf{x}_i(1) = \mathbf{T}^{-1}\mathbf{x}_i(1) + \mathbf{v}$ and the inverse of the masses with not changing the expected frequencies. The vector $\mathbf{v}$ changes the mean position of the row points on each dimension whereas the diagonal matrix $\mathbf{T}$ changes the spread of the row points on each dimension. These transformations, however, do not change the order or relative spacings between row points; they are a classic transformation. Similar transformations can be applied to the column points and masses, by using the inverse of this linear transformation. To obtain optimal location ($\mathbf{a}$) and scalings ($\mathbf{T}$) the correlation between squared distances $(d_{ij}^2(\mathbf{Z}_1, \mathbf{Z}_2))$ and $\hat{F}_{ij}/\hat{m}_i\hat{m}_j$ is minimized. This can be done by using the program PROFIT (Commandeur, 1991).

It is important that the total mass is equal over the time points, i.e. that the total mass at the first time point is equal to the total mass at the second time point. Therefore the identification constraints on the masses were also such that this restriction is true. The mass will be represented by the axes of the circle. To these the circle we shall therefore use a radius equal to $r(i) = \sqrt{|m(i)|/\pi}$.

3.7. Estimation

Several researchers have discussed estimation of the RC(*M*) association model (Goodman,

two-way tables, e.g. transition data that are obtained in different countries, or different groups or at different time points.

4.1. The model

The gravity models that shall develop models for multiple two-way tables. Each of these tables can be modelled by the gravity models of the previous section, resulting in a universe with objects and masses for each layer ($k = 1, \ldots, K$) of the table. The most general model is

$$F_{ijk} = \frac{m_{ik}(\alpha m_{jk})}{\exp(\tau_{ik}^2 / 2\sigma_{ik}^2 \cdot \mathbf{Z}_{1k} \cdot \mathbf{Z}_{2k})}, \tag{9}$$

where $m_{ik}(k)$ is the mass of object i at time point i in layer k and the vector $\mathbf{z}_i(k)$ gives the position of object i at time point i in layer k, i.e. each layer is represented by the gravity model for one two-way table.

Restrictions can be imposed to relate the different universes. For example, the masses of different layers or the co-ordinates of different layers can be constrained to be equal or equal up to a multiplicative constant. The most natural choice is to restrict the co-ordinates ($\mathbf{Z}_{1k}$). Examples of restrictions are

$$\mathbf{Z}_{1k} = \mathbf{Z}_1 \mathbf{W}_k, \tag{10}$$

$$\mathbf{Z}_{1k} = \mathbf{Z}\mathbf{W}_k, \tag{11}$$

$$\mathbf{Z}_{1k} = \mathbf{Z}_1, \tag{12}$$

$$\mathbf{Z}_{1k} = \mathbf{Z}, \tag{13}$$

where $\mathbf{W}_k$ is a diagonal matrix, specifying positive weights that stretch or shrink the dimensions of $\mathbf{Z}$ or $\mathbf{Z}_1$ and $\mathbf{Z}$ is a matrix containing the row points of the first occasion and the column points ($\mathbf{Z}_1$) over $(\mathbf{Z})$. When $r_{\mathbf{W}_k} < 1$, dimensions of the layer k is restricted, meaning that the layer k the objects are more differentiated on this dimension. When $r_{\mathbf{W}_k} = 1$ the dimension shrinks, i.e. for layer k the objects are less differentiated on this dimension. Restriction (10) corresponds to the well-known INDSCAL type of three-way distance models (Carroll and Chang, 1970), with stable positions of the categories for each layer of the table. The first restriction, equation (10), is the most general restriction. Restriction (11) specifies the model with stable but unrelated dynamic positions. The third restriction, equation (12), follows two-mode distances which are equal across the layers and the fourth restriction, equation (13), defines one-mode distances that are equal across the layers. An example of a model with stable positions within each layer (but is stretched or shrunk in the different layers and with equal masses over the layers can be found in de Rooij (2001).

4.2. Rewriting the model

Like the models for two-way tables this model can be written as an association model. In this case we deal with the conditional association models as proposed by Clogg (1982) and Becker and Clogg (1989). Although this is not necessary, it is convenient to use the available computer software for fitting our gravity models and helps in understanding relationships between our gravity models and other models for contingency tables. The formulae for transforming one model into the other are similar to expressions (6) and (7). For example, with the restriction in equation (10) we have

$$\begin{aligned} F_{ijk} &= \frac{m_{ik}(\alpha m_{jk})}{\exp(\tau_{ik}^2 / 2\sigma_{ik}^2 \cdot \mathbf{Z}_1 \mathbf{W}_k \cdot \mathbf{Z}_2 \mathbf{W}_k)} \\ &= \frac{m_{ik}(\alpha m_{jk})}{\exp(\tau_{ik}^2 / 2\sigma_{ik}^2) \exp(-\tau_{ik}^2 / 2\sigma_{ik}^2 \cdot \mathbf{Z}_1 \mathbf{W}_k \cdot \mathbf{Z}_2 \mathbf{W}_k)} \\ &= \frac{m_{ik}(\alpha m_{jk})}{\exp(\tau_{ik}^2 / 2\sigma_{ik}^2) \exp(\tau_{ik}^2 / 2\sigma_{ik}^2 \cdot \mathbf{Z}_1 \mathbf{W}_k \cdot \mathbf{Z}_2 \mathbf{W}_k)} \\ &= \alpha_{ik}\beta_{jk} \exp\left(\sum_{s=1}^{S} \phi_{sk} x_{is}x_{js}\right), \end{aligned} \tag{14}$$

where the difference between α_{ik} and α_{ik} is a scaling factor, β_{jk} and $\phi_{sk} = \phi_s$. The last line is expression (14) in the conditional association model (Clogg, 1982; Becker and Clogg, 1989).

The gravity models can be rewritten in a form that is well known for two-way table. By using equation (14) (15) the co-ordinates of the different layers can be obtained, but with the same restrictions (11) and (13) symmetry restrictions on the association are imposed, whereas without

from (11) and (13) result in models without three-way association.

4.3. Identification

The conditional association model needs location and scaling constraints but on cross-dimensional constraints (Clogg, 1982). Since the RC(*M*) association model is identified, as discussed in the previous section, new locations ($\mathbf{a}$) and scalings ($\mathbf{T}$) can found by minimizing the correlation between the elements α_{ik} defined as

$$\alpha_{ik} = \frac{\alpha_{ik}}{1 + \frac{\alpha_{ik}}{\exp \mathbf{a}_k}}$$

and the squared unweighted two-mode distances. The one-mode distance models (11) and (13) are identified.

The number of freedom for the conditional association model were discussed in Wong (2001), pages 204–207. For our models these numbers should be adapted for the loyalty parameters for the non-movers, i.e. the off frequencies represented within parentheses in Table 2. The model that is given in expression (9) is multiplied by the term $\exp(\tau_{ik}\lambda_{ik})$ where λ_{ik} equals 1 if $i = j$ and 0 otherwise. There are IK of these parameters.

4.4. Estimation

The conditional association models can be estimated with LEM (Vermunt, 1997). From the expected frequencies we obtain an identified solution with a MATLAB procedure.

A cautionary note is in order here: when the model with constraint (10) or (11) is estimated with LEM, sometimes negative estimates for the association parameters are obtained. On the contrary level, negative association parameters may occur for some layer. In that case there is not a distance representation of the conditional association model. A traditional (i.e. for the association model) graphical display of the association may also reflect the dimension for the row or column points for the specific layer. If such a negative association coefficient occurs it is

Table 5. Mobility data from three countries (USA, UK and Japan)

Country	Father	Son				
		$\frac{\text{WC}}{\text{WC}}$	$\frac{\text{WC}}{\text{BC}}$	$\frac{\text{BC}}{\text{WC}}$	$\frac{\text{BC}}{\text{BC}}$	$\frac{\text{Sum}}{\text{Sum}}$
USA	WC	1414	521	724	256	2915
	WC	1807	1077	759	445	4 088
	BC	1409	790	1318	1179	4 696
	BC	1197	791	1775	2790	6 553
UK	WC	1855	1004	1	1	2 861
	WC	4758	1778	17	7	6 560
	BC	1	1	1	1	4
	BC	2	1	1	1	5
Japan	WC	1324	1	1	1	1 327
	WC	2011	1	1	1	2 015
	BC	1	1	1	1	4
	BC	1	1	1	1	4

Of male, Yamaguchi (1979, p. 75, gathers men scoring 0, 1, lower non-manual, 2 lower manual, 3 upper manual, 4 upper non-manual. Of female, Yamaguchi (1979, p. 75, gathers women scoring 0, 1, lower non-manual, 2 lower manual, 3 upper manual, 4 upper non-manual.

The generalization to tables for three time points needed a further expression of the law of gravity. Distances between three points, triadic distances, were discussed and their relationship to partial association models was shown. Again the triadic distance is easier to understand than the sum of three inner products, as modified in the partial association models. Although this generalization is not unique, it appears to be the only one that is based on the concepts of *mass* and *distance*. The models that were discussed can theoretically be generalized to more time points by using tetradic (or polyadic) distances, which can be defined as sums of all (squared) bivariate, triadic, tetradic, ..., $(I-1)$-adic distances. The relations between the gravitational models and triadic distance models as shown in equations (13) and (25) does not generalize to more time points (variables), for that case the gravity models become more restricted than the association models.

The triadic distance models are good at showing and representing change. They look a formal analysis of change in the sense that they are invariant under the group of rotations. The triadic gravity models can be *scanned* as second-order Markov models with restrictions, and *when* interpreted by using the correct temporal ordering there is an influence of the first time point on the third time point, mediated by the second time point, as discussed in Section 3.1. For a detailed examples of interpretation at the end of Section 3.1. For triadic three-mode distances similar statements about conditional odds can be obtained.

The triadic distance model is different from the change-from-time-point-1 to-2 in triadic distance models is different from the change-that is obtained when the table was collapsed over the final time point. A model is *collapsible* when the conditional association equals the marginal association, and this is generally so for conditional independence models (Bishop *et al.*, 1975). For the example that was discussed in Section 3 this would be the non-fitting first order Markov model. Collapsibility is discussed in the context of causal modeling and path analysis, but also in ignoring relevant covariates in the analysis (Agresti (2002), page 478). In other words, the group of people who make the same change from time point 1 to 2 do not form a homogeneous group by using the *distance* that are made at the third time point we obtain more reliable change scores.

In this paper all models were estimated with LEM. This yielded some problems; for example, the triadic distance models for the Dutch mobility tables could not be estimated with the available software for association models. Furthermore, for models for observations on three times using ordered categories the number of cells in the table can become too large. Finally, the models for multiple two-way tables sometimes a negative association coefficient occurs for a particular layer which under a distance model is not possible. To deal with such problems special software should be written.

We tried to build models for the whole data set. Another way of analyzing square tables is to decompose the table into a symmetric part and a skew symmetric part, and these are then further decomposed (Bishop *et al.*, 1975). An interesting property of one of the symmetry decompositions is that due to Gower (1977) and discussed also in Constantine and Gower (1978). Bilinear forms for skew symmetry were discussed in van der Heijden and Mooijaart (1995). For a treatment of skew symmetry in the three-way case see de Ruiij and Bijmat (1998).

The use of gravity models to explain social phenomena is not new. Tobler (1976), for example, discussed gravity models for migration and spatial interaction. Applications of gravity models in economics and transportation studies (see, for example, Sen and Smith (1995)). In these cases, however, the distances are often known in advance, *i.e.* they are real distances, or the distance between two points is measured in time. Then the gravity model is used to explain the variation in the data. The fact that Newton's gravity model is used more often in other areas is due to its simple and understandable nature: mass and distance are easily grasped concepts. In this paper

these concepts are used for the analysis of change. For change from one time point to another these models give a very natural description of the change process.

Acknowledgements

This project was finished while the author was sponsored by the Netherlands Organization for Scientific Research, international grant 47-516-002. I thank Paul Eilers, Willem Heiser, Carolyn Anderson and the referees for discussions and comments on earlier versions of the manuscript.

Appendix A: Software note

We used the program LEM to obtain fitted frequencies. The scaling and locations were found by using MATLAB. A MATLAB-shell was built around LEM and the m-manual copying is needed to perform the analysis. On request the MATLAB and LEM files can be obtained from the author.

References

Agresti, A. (2002) *Categorical Data Analysis*, 2nd edn. New York: Wiley.
Anderson, C. J. and Vermunt, J. K. (2000) Log-multiplicative association models as latent variable models for nominal and ordinal data. *Sociol. Methodol.* **30**, 81–121.
Becker, M. P. (1994) Algorithm AS 291: Maximum likelihood estimation of the RC(*M*) association model. *Appl. Statist.* **39**, 152–167.
Becker, M. P. and Clogg, C. C. (1989) Analysis of sets of two-way contingency tables using association models. *J. Am. Statist. Ass.* **84**, 142–151.
Bishop, Y. M. M., Fienberg, S. E. and Holland, P. W. (1975) *Discrete Multivariate Analysis: Theory and Practice*. Massachusetts: Massachusetts Institute of Technology Press.
Bock, R. D. (1975) *Multivariate Statistical Methods in Behavioral Research*. New York: McGraw-Hill.
Carroll, J. D. and Chang, J. J. (1970) Analysis of individual differences in multidimensional scaling via an N-way generalization of "Eckart-Young" decomposition. *Psychometrika* **35**, 283–319.
Constantine, [G.] (1985) Contributions to Canadian participation in the gold rush. II. Contributions to the statistical analysis of contingency tables. *Can. Soc. Sci. Forum* **28**, 701–726.
Constantine, [G.] and Gower, J. C. (1978) Graphical representation of asymmetry. *Appl. Statist.* **27**, 297–304.
Coxon, A. P. M. (1982) *The User's Guide to Multidimensional Scaling*. London: Heinemann.
Gabriel, K. R. (1971) The biplot graphic display of matrices with application to principal component analysis. *Biometrika* **58**, 453–467.
Gilula, Z. and Haberman, S. J. (1986) Canonical analysis of contingency tables by maximum likelihood. *J. Am. Statist. Ass.* **81**, 780–788.
Gilula, Z. and Haberman, S. J. (1988) The analysis of multivariate contingency tables by restricted canonical and restricted association models. *J. Am. Statist. Ass.* **83**, 760–771.
Gilula, Z. and Krieger, A. M. (1989) The decomposition of categorical data: an approach and an application. *Can. J. Statist.* **17**, 59–74.
Gilula, Z. and Ritov, Y. (1990) Inferential ordinal correspondence analysis: motivation, derivation and limitations. *Int. Statist. Rev.* **58**, 99–108.
Gilula, Z. and Ritov, Y. (1991) Canonical analysis of contingency tables by maximum likelihood: theory and applications. *Ann. Statist.* **19**, 1538–1562.
Gilula, Z. and Ritov, Y. (1992) Asymptotic normality of canonical correlation analysis based on contingency tables by maximum likelihood. *J. Multivar. Anal.* **41**, 1–14.
Gilula, Z. and Ritov, Y. (1994) Inference in ordinal correspondence analysis: motivation, derivation and limitations. *J. Math. Psychol.* **38**, 307–358.
Gilula, Z. and Ritov, Y. (1995) Canonical analysis of two-way contingency tables by maximum likelihood: theory and applications. *Can. J. Statist.* **23**, 135–147